

# MAPPING OF HEAVY METAL CONTAMINATION IN ALLUVIAL SOILS OF THE MIDDLE NILE DELTA OF EGYPT

Mohamed S. Shokr<sup>a,b</sup>, Ahmed A. El Baroudy<sup>a</sup>, Michael A. Fullen<sup>b</sup>, Talaat R. El-beshbeshy<sup>a</sup>, Rafat R. Ali<sup>c</sup>, Abd El Halim A. Abd El Halim<sup>a</sup>, Antonio J.T. Guerra<sup>d</sup> and Maria C.O. Jorge<sup>d</sup>

<sup>a</sup>*Soils and Water Department, Faculty of Agriculture, Tanta University, Tanta, Egypt.*

<sup>b</sup>*The University of Wolverhampton, Wolverhampton WV1 1LY, UK.*

<sup>c</sup>*Soils and Water Use Department, National Research Centre, Giza, Egypt.*

<sup>d</sup>*Department of Geography, Federal University of Rio de Janeiro, Brazil.*

*E-mail: m.s012@yahoo.com (corresponding author: Mohamed Shokr)*

Submitted and accepted dates (use style Received dates, or Alt + Ctrl + R)

**Abstract.** Areas contaminated by heavy metals were identified in the El-Gharbia Governorate (District) of Egypt. Identification used remote sensing and Geographical Information Systems (GIS) as the main research tools. Digital Elevation Models (DEM), Landsat 8 and contour maps were used to map physiographic units. Nine soil profiles were sampled in different physiographic units in the study area. Geochemical analysis of the 33 soil samples was conducted using X-ray fluorescence spectrometry (XRF). Vanadium (V), nickel (Ni), chromium (Cr), copper (Cu) and zinc (Zn) concentrations were measured. V, Ni and Cr concentrations exceeded recommended safety values in all horizons of the soil profiles, while Cu had a variable distribution. Zn concentrations slightly exceeded recommended concentration limits. Concentrations were mapped in each physiographic unit using the inverse distance weighted (IDW) function of Arc-GIS 10.1 software. Pollution levels were closely associated with industry and urban areas.

**Keywords:** Soil contamination, X-ray fluorescence spectrometry, remote sensing, Geographical Information Systems, Middle Nile Delta, Egypt.

## Introduction

Sustainable agriculture is mainly related to environmental, agronomic, ethical and socio-economic issues (Abd Elgawad et al. 2007). One aspect of sustainability is accumulation of heavy metals in soils, which may cause serious problems, if certain levels are exceeded. In recent years, much concern has been articulated over problems of soil contamination with heavy metals. These metals can accumulate in plants and animals and then in humans through the food chain (Govil et al. 2001; Lu, Bai; 2010; Romic, Romic, 2003). Thus, heavy metals may damage human health and the environment (Jankaite, Vasarevičius, 2005).

The Nile Delta (area ~20,000 km<sup>2</sup>) represents only 2.3% of the area of Egypt, but it has ~46% of the total cultivated area (55,040 km<sup>2</sup>) and accommodates ~45% of Egypt's population (Fanous 2002), with densities ≤1600 inhabitants per km<sup>2</sup> (Zeydan 2005). On the Nile Delta ~63% land is agricultural, due to suitable soil properties and the presence of irrigation systems (Dawoud 2004). The River Nile divides into two branches, the Rosetta and Damietta, and the Delta region is located between them (Dumont, 2009). The Nile Delta (area 404,686 ha) depends on drainage water for irrigation (Abu Khatita 2011).

There are three major layers in the middle Delta aquifer (Atwia et al. 2006). The uppermost layer is composed of clay deposits, the second layer is formed from sandy clay deposits and the third layer is composed of saturated sand and gravel. Thus, the thin clay layers and presence of sandy clay lenses facilitate percolation of sewage water to the aquifer. Many activities, including agricultural development and industrial activities and inadequate rural sanitation, have impacts on eutrophication and contamination status, ecological value and environmental conditions in the Nile Delta (Zeydan 2005).

Heavy metal contamination of soil may present risks and hazards to humans and the ecosystems through: direct consumption or contact with contaminated soil, the food chain (soil-plant-human or soil-plant-animal-human), drinking of contaminated ground-water, decreased food quality (safety and marketability) via phytotoxicity, reduction in land usability for agricultural production causing food insecurity, and land tenure problems (McLaughlin et al. 2000; Ling et al. 2007). Huge amounts of fertilizers are regularly added to soils in intensive farming systems to provide sufficient nitrogen (N), phosphorus (P) and potassium (K) for crop growth. The compounds used to supply these elements contain trace amounts of heavy metals which, after continued fertilizer application, may significantly increase soil metal contents (Raven et al. 1998). Integration of remote sensing information within a GIS database can quickly provide detailed soil survey information at low cost. GIS databases can also help derive Digital Elevation Models (DEM), which can help derive landscape attributes utilized in landform characterization (Brough 1986, Dobos 2000).

It is critical to analyse the distribution and concentration of metals. This will enable identification of contamination levels and assess associated impacts, on both the environment and human health. Soils are a vital sink for these metals,

because of their high metal retention capacities. The assessment and mapping of soil heavy metals can assist the development of strategies to promote sustainable use of soil resources, decrease soil degradation and expand crop production. Remediation of soils polluted by heavy metals is a major global ecological issue. Remote sensing is one of the most important methods used for soil survey, mapping and environmental investigations (Lillesand and Kiefer 2003). Geostatistical interpolation is used to survey and interpret the spatial distribution of pollutants in soil (He, Jia 2004; Woo et al., 2009). The inverse distance weighted (IDW) function is helpful when the purpose is to investigate overall pollution patterns (Zheng 2006). The Middle Nile Delta is affected by different pollution sources, because of the increasing number and types of industries, urban expansion, increased traffic volumes, use of drain-water and waste deposits (Abu Khatita 2011). The latter may well present a long-term danger. Usually waste deposits just settle within the normal Nile sediments and no special effort is made to construct barriers, which hinder the migration of water from these deposits into ground-water. High concentrations of vanadium (V) can damage human health, while the inhalation of airborne V-compounds can affect eyes, throat and lungs, produce weakness, ringing in the ears, nausea, vomiting, headaches and damage nerve systems (Lagerkvist, Oskarsson 2007). In Egypt, measured chromium (Cr) contents in soils range between 11.6-179 ppm, and depend on soil types and land management (Abdel-Sabour et al. 2002). Cr toxicity depends on its oxidation status. While  $\text{Cr}^{3+}$  is considered relatively harmless,  $\text{Cr}^{6+}$  is highly toxic. Cr uptake can cause diarrhoea, bleeding in the stomach and intestines, liver and kidney damage and cramp. Nickel (Ni) compounds are relatively non-toxic for plants and animals, but there is an increased risk of respiratory tract cancer, due to exposure to nickel sulphide and oxides (Sundermann, Oskarsson, 1991).

Copper (Cu) is an essential element for all life-forms. In plants, Cu is required in small amounts (5-15 ppm) (Bowen 1979). The amount of Cu in soils may affect crop growth and yields. The application of Cu salts to Cu-deficient soils increases crop yields, because it compensates for Cu deficiency in plants (Baker, Senft, 1995). Coal fly-ash contains 48  $\mu\text{g/g}$  of Cu (Wong, Wong, 1986). In Ohio (USA), measured Cu concentrations in indoor dust were twice that of outdoor dust (Tong 1998). Cu toxicity in humans is relatively rare, because they can tolerate levels  $\leq 12$  mg/day (WHO 1996). However, Cu deficiency in humans causes anaemia, bone and cardiovascular disorders, mental and nervous system deterioration and defective keratinization of hair.

Zinc (Zn) is the fourth most used metal in the world, after iron (Fe), aluminium (Al) and Cu (Bradl, Xenidis, 2005). Zn uptake can lead to health disorders, including pancreatic diseases. Inhalation of Zn-oxide (particle size 0.2-1  $\mu\text{m}$ ) during Zn-processing causes metal fume fever, which is characterized by a sore throat, cough, fever, vomiting and pneumonitis (Ohnesorge, Wilhelm, 1991).

The main aim of this research is to identify land contaminated by heavy metals in the El-Gharbia Governorate (District) of Egypt. This was undertaken using remote sensing, Geographical Information Systems (GIS) and X-ray fluorescence (XRF) spectrometry (Fig. 1).




Fig. 1. Location of the study area in the middle Nile Delta of Egypt

 Study area

## 1. Materials and methods

### 1.1. Study area

The study area occupies the Middle part of the Nile Delta of Egypt. It is bounded by 30°45'20"-31°10'50"E and 30°35'10"-31°10'05"N, and covers an area of 1927.4  $\text{km}^2$  (Fig. 2). Based on the US Soil Taxonomy (USDA 2010) the soil temperature regime of the study area is Thermic and the soil moisture regime is Torric. The mean annual

temperature reaches its maximum in June, July and August and often exceeds 30°C. The mean minimum temperature (11.2°C) usually occurs in January, February or March at Tanta Meteorological Station (Climatologically Normal for Egypt 2011). Precipitation is unequally distributed through the rainy season. Annual rainfall is very low and mostly falls in winter; with a mean 3.8 mm/year. Rain mainly falls in the cold season (November-March) and the minimum amount is in June and September. The area belongs to the late Pleistocene era, which is evidenced by the deposits of the Neogene, which are composed of medium and fine silts (Said 1993).




Fig. 2. Landsat 8 mosaic of the study area

## 1.2. Digital image processing and physiographic mapping

Digital image processing was completed for two Landsat 8 satellite images (path 177/row 38 and path 177/row 39), with a spatial resolution of 30 m, acquired in May 2014. The images were pre-processed, including radiometric correction (used to modify digital values of pixels to remove noise). Images were geometrically rectified using the Universal Transverse Mercator (UTM) co-ordinates, with the World Geodetic System datum (WGS 1984) and then maps were constructed. Images were atmospherically corrected using the FLAASH module (ITT 2009). Data were calibrated to radiance using the inputs of image type, acquisition date and time. Images were subject to linear stretching by 2%, smooth-filtered, and their histograms were matched, adopting the procedures of Lillesand and Kiefer (2007) and mosaicked using ENVI 5.1 software. The extraction of landform units used high spatial resolution images, so the spatial resolution of satellite image was enhanced using the data merge function of Envi5.1 software. Merging is performed by using multispectral bands (~30 m) as low spatial resolution, and band 8 (panchromatic band) with ~15 m resolution. Landforms were extracted using contour maps (scale 1:25,000) and enhanced satellite images. Both enhanced satellite images were processed with DEM in ERDAS Imagine 8.7, to extract the landform information (Dobos et al. 2002). The initial landform maps were ground-truthed using field observations.

## 1.3. Spatial distribution of heavy metals

Spatial interpolation is widely used when data are collected at distinct locations (e.g. soil profiles) for producing continuous information (Ali, Moghanm 2013). Inverse distance weighted (IDW) is an interpolation method which uses measured values surrounding the prediction location. The measured values closest to the prediction location have more influence on the predicted value than those farther away, thus giving greater weight to points closest to the prediction location, and the weights decrease as a function of distance (Shepard 1968). Geostatistical relationships among the known points (IDW) of Arc-GIS 10.1 software were used to interpolate heavy metal concentrations in the study area. The spatial interpolation method (IDW) was used with 12 neighbouring samples for estimation of each grid point. A power of two was used to weight the nearest points.

#### 1.4. Assessment of contamination risk

The Geoaccumulation Index (I<sub>geo</sub>) was originally used to evaluate bottom sediment contamination. However, it has been successfully used to evaluate soil contamination (Gowd et al. 2010). The I<sub>geo</sub> Index means the assessment of contamination depends on comparing heavy metal concentrations in soils to background values. The calculation of the Geoaccumulation Index uses the equation:

$$I_{geo} = \log_2 \frac{C_n}{1.5B_n} \quad (1)$$

Where C<sub>n</sub> = the measured concentration of the element in soil.

B<sub>n</sub> = the geochemical background concentration of the heavy metal.

The Geoaccumulation Index (I<sub>geo</sub>) is shown in Table 1.

Table1. The Geoaccumulation Index (I<sub>geo</sub>) for assessing contamination levels in soil (source: Rahman et al. 2012)

I geo Class	I geo value	Contamination level
0	$I_{geo} \leq 0$	Uncontaminated
1	$0 < I_{geo} < 1$	Uncontaminated/moderately contaminated
2	$1 < I_{geo} < 2$	Moderately contaminated
3	$2 < I_{geo} < 3$	Moderately/strongly contaminated
4	$3 < I_{geo} < 4$	Strongly contaminated
5	$4 < I_{geo} < 5$	Strongly/extremely contaminated
6	$5 < I_{geo}$	Extremely contaminated

#### 1.5. Soil analysis

Soil samples were collected from nine profiles in El-Gharbia Governorate. The selected profiles represent the different soil units. Pedological descriptions of profiles were conducted using the procedures of FAO (2006) (Table 2). About 1 kg was collected from each horizon of each profile. Soil samples were air-dried and large stones and organic debris were removed before sieving. Samples were gently ground, homogenized, sieved through a 2.0 mm sieve and then crushed to a fine (<125 µm) powder. Oven-dry samples were ignited at 375°C for 16 hours (overnight), adopting the procedures of Ball (1964). Subsamples of 8.5 g of soil powder were added to 1.5 g of wax (Lico waxc micropowder PM, Hoechst wax) and then compressed under 12 tonnes pressure by a semi-automatic hydraulic press to make a pellet. The geochemical composition of soil pellets were analysed using an XRF spectrometer model Epsilon3 XLE. XRF analyses were performed at the University of Wolverhampton, UK.

Table 2. Pedological descriptions of soil profiles

Profile Number	Depth (cm)	Colour	Texture	Structure	Soil consistency
1	0-50	5YR 3/4 5YR5/4	Loam	Sub-angular blocky	Sticky, plastic
	50-85	7.5YR 5/6 7.5YR 6/6	Sandy loam	Sub-angular blocky	Slightly sticky, slightly plastic
	85-120	10YR 5/8 10YR 6/8	Loam	Sub-angular blocky	Sticky, plastic
	120-150	10YR 5/8 10YR 7/8	Sandy loam	Sub-angular blocky	Slightly sticky, slightly plastic
2	0-45	5YR 4/6 5YR 5/6	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	45-85	10YR 5/8 10YR 6/8	Sandy Clay Loam	Sub-angular blocky	Sticky, plastic
	85-110	10YR 5/8 10YR 7/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
3	0-75	10 YR 5/6 10YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	75-100	10YR 5/6 10YR 7/8	Loam	Sub-angular blocky	Sticky, plastic
	100-150	10YR 5/8 10YR 7/8	Silt loam	Sub-angular blocky	Sticky, plastic
4	0-60	5YR 4/8 5YR 4/6	Loam	Sub-angular blocky	Sticky, plastic
	60-100	10YR 5/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	100-120	10YR 7/6	Loam	Sub-angular blocky	Sticky, plastic
	120-150	5YR 4/8 5YR 4/6	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
5	0-45	10YR 5/8 10YR 7/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	45-65	7.5YR 5/6 7.5YR 5/8	Loam	Sub-angular blocky	Sticky, plastic
	65-110	10YR 5/8 10YR 6/8	Silt loam	Sub-angular blocky	Sticky, plastic
	110-150	10YR 5/8 10YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
6	0-35	10YR 5/6 10YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	35-65	10YR 6/8 10YR 7/8	Loam	Sub-angular blocky	Sticky, plastic
	65-100	10YR 5/8 10YR 7/8	Loam	Sub-angular blocky	Sticky, plastic
	100-150	5YR 5/6 5YR 4/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
7	0-55	5YR 4/8 5YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	55-110	5YR 4/8 5YR 6/8	Loam	Sub-angular blocky	Sticky, plastic
	110-150	7.5YR 5/8 7.5YR 6/4	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
8	0-30	7.5YR 4/4 7.5YR 5/8	Silt loam	Sub-angular blocky	Sticky, plastic
	30-60	7.5YR 4/6 7.5YR 6/8	Silt loam	Sub-angular blocky	Sticky, plastic
	60-100	7.5YR 5/8 7.5YR 6/8	Silt loam	Sub-angular blocky	Sticky, plastic
	100-150	7.5YR 5/8 7.5YR 5/8	Silt loam	Sub-angular blocky	Sticky, plastic
9	0-45	7.5YR 5/8 7.5YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	45-105	10YR 5/8 10YR 6/8	Loam	Sub-angular blocky	Sticky, plastic
	105- 130	10YR 5/8 10YR 6/8	Sandy Loam	Sub-angular blocky	Slightly sticky, slightly plastic
	130-150	7.5YR 6/8 7.5YR 5/8	Loam	Sub-angular blocky	Sticky, plastic

## 2. Results and Discussion

### 2.1. Physiographic map of the study area

The satellite images show that the study area is a flood-plain and includes high terraces (12.04% of area), moderately high terraces (22.41%), low terraces (21.67%), high decantation basins (2.05%), low decantation basins (12.26%), high overflow basins (12.68%), low overflow basins (10.71%), river levees (5.38%) and swales (0.77%). The main physiographic soil units of the study area are reported in Table 3 and Fig. 3.

Table 3. Physiographic units on the soil map

Physiographic unit	Landforms	Mapping unit	Soil profile	Profile elevation (masl)	Area (km <sup>2</sup> )	Area (% )
	High terraces	T1	9	12	232.21	12.05
	Moderately high terraces	T2	4	8	431.99	22.41
	Low Terraces	T3	1	0	417.8	21.68
	High Decantation Basin	D1	3	10	39.53	2.05
Flood plain	Low Decantation Basin	D2	5	6	236.29	12.26
	High overflow Basin	OB1	6	7	244.461	12.68
	Low Overflow basin	OB2	7	5	206.45	10.71
	Levees	L	8	9	103.82	5.39
	Swales	S	2	8	14.89	0.77
	Total	-	-	-	1927.441	100.00

\*masl = metres above sea level.

1: River terraces: these soils represent the late Pleistocene deltaic plain and occur at the edge of decantation basins (these are basins in which sedimentation, particularly of silt and clay, occur during floods). The soils are formed on terraces at various heights above the valley floor.

2: Basins: these are artificially enclosed areas of a river or harbour, designed so that water levels are unaffected by tides.

3: River levees: these are a type of dam that runs along the banks of rivers or canals. Levees reinforce the banks and help prevent flooding. By confining the flow, levees can also increase water velocity.

4: Swales: these are low tracts of land, usually consisting of moist and marshy lands. The term can refer to both natural and artificial landscape features. Artificial swales are often designed to manage water runoff, filter pollutants and increase rainwater infiltration.




Fig. 3. The main landforms of the study area and profile locations

## 2.2. Heavy metal contamination

XRF analyses of the soil samples identified the presence of  $\text{SiO}_2$ ,  $\text{Al}_2\text{O}_3$ ,  $\text{P}_2\text{O}_5$ ,  $\text{K}_2\text{O}$ ,  $\text{CaO}$ ,  $\text{MgO}$ ,  $\text{Na}_2\text{O}$  and  $\text{Fe}_2\text{O}_3$  (major) and Cr, Cu, Zn, Ni, Br, Rb, Sr, Y, Zr, Nb, Sn, Te, Ba, Eu, Yb, Re, Ga, Ir, Mo, As and Pb (minor). Concentrations of the heavy metals Cr, Cu, Ni, V and Zn for each profile are reported in Table 4. For the metals Te, Mo, As and Pb, results are not reported, because their concentrations were below detection limits. Spatial interpolation maps (Figs. 4, 5, 6 and 7) of heavy metal concentrations were prepared using the IDW function (inverse distance weighted) interpolation method in Arc GIS 10.1.

### Vanadium

The concentrations and the interpolation map for V in the soil samples are given in Table 4 and Fig. 4. V concentrations ranged from 194.0-744.4 mg/kg with a weighted mean ranging from 206.79-450.58 mg/kg (Table 5). The highest measured concentration of V was in the upper horizon of Profile 2, which represents a swales unit and is located 270 m north of Mansuriyyat Al-Farastaq village, ~6.5 km south-west from the centre of the town of Kfr Elzayat (population in 2015 was 448,965). The Igeo Index showed that all soil samples were in the uncontaminated/moderately contaminated categories, except for first horizon of Profile 2 in the swales mapping unit, which is classified as moderately/strongly contaminated (Table 6). The high deposition of V might be caused by the numerous local factories. V concentrations are higher than the permissible limits (90 mg/kg), recommended by Bowen (1979) in all soil profile horizons (Table 5). The spatial interpolation shows an increasing trend from north-east to south-west. The highest weighted mean (weighting concentration by representative area) (450.58 mg/kg) was found in 0.77% of the study area. From the interpolation map of V in the study area (Fig. 4) we can conclude that the order of concentration ascending in the mapping units is: low decantation basin (D2), high overflow basin (OB1), low overflow basin (OB2), moderately high terraces (T2), high decantation basin (D1), levees (L), high terraces (T1), low terraces (T3) and swales (S).




Fig. 4. Spatial interpolation of the weighted mean of vanadium

## Chromium

Anthropogenic sources of Cr include alloys, chrome plating, pigments, chemical catalysts, dyes, tanning, wood impregnation and refractory bricks (Reimann, de Caritat, 1998). The highest concentration of Cr (519 mg/kg) was in the top-soil of Profile 2, which may be due to the many local factories. The lowest (140.3 mg/kg) was in the top-soil of Profile 6, which represents a high over-flow basin (Table 4). The mean weight of Cr concentrations ranged from 152.84-314.73 mg/kg (Table 5). All concentrations exceeded the recommended values given by Bowen (1979) and, according to the Igeo Index, most soil samples are in the uncontaminated/moderately contaminated category (Table 6). Cr concentrations increased from east to west and south of the study area (Fig. 5). The highest Cr concentrations tended to be in the swales unit and the lowest in the high overflow basin unit.






Fig. 5. Spatial interpolation of the weighted mean of chromium

#### Nickel

Baghdady, Sippola (1984) reported that the mean total Ni content in Egyptian alluvial soils is 64.4 mg/kg, ranging from 20-74 mg/kg, while the mean  $\text{NH}_4\text{OAC-EDTA}$  extractable Ni is 1.9 mg/kg, ranging from 1.0-2.2 mg/kg. However, in this study, Ni concentrations ranged from 60.60-267.30 mg/kg (Table 4), with mean weight ranging from 69.60-148.39 mg/kg (Table 5). Ni concentrations were higher in alluvial soils than previous studies and exceeded the permissible limit (50 mg/kg) (Bowen 1979). According to the Igeo Index, all samples are in the uncontaminated and moderately contaminated categories (Table 6). Table 4 reports Ni concentrations in soil profiles and Fig. 6 shows the spatial trends, which increased from north to south and west. The highest concentrations were in swales, which occupy 14.89 km<sup>2</sup> of the study area. The interpolation of Ni shows high spatial variability, with the lowest values in the high decantation basin units.




Fig. 6. Spatial interpolation of the weighted mean of nickel

#### Copper

Table 4 and Fig. 7 report Cu concentrations and the spatial interpolation of weighted mean Cu concentrations, respectively. Cu contents in horizons ranged from 0-288.9 mg/kg and mean weight ranged from 28.90-152.92 mg/kg (Table 5). All concentrations exceeded the permissible limit of 25 mg/kg (Bowen 1979), except for the second horizon of Profile 4 and the first, second and the third horizons of Profile 8 (Table 4). In addition, in Profile 9 the deepest horizon exceeded the limit, whereas concentrations in the upper layer were 0. This is probably due to percolation and illuviation of Cu associated with irrigation water. These profiles represent moderately high terraces, levees and high terraces, respectively. The Igeo Index showed that soil samples were in three contamination categories (uncontaminated/moderately contaminated, moderately contaminated and moderately/strongly contaminated) (Table 6). Two important sources of Cu in the Nile Delta are: (i) applications of Cu-based liquid fungicides, and (ii) use of  $\text{CuSO}_4$  as an algicide in treating and controlling problematic macro-algal blooms in the Nile, especially during summer (Abdel-Moati, El-Sammak 1997). The lowest Cu concentrations were in the river levees units and the highest values were in the swale units.




Fig. 7. Spatial interpolation of the weighted mean of copper

#### Zinc

Zn concentrations slightly exceed the permissible concentration limit of 90 mg/kg (Bowen 1979) (Table 5). Exceptions include the upper layers of Profiles 2 and 4, where concentrations greatly exceeded permissible limits (377.6 and 308.0 mg/kg, respectively). The highest concentrations were in the upper horizon, but in Profile 8 the highest concentration was in the subsurface (Table 4). This could be caused by infiltration of irrigation water through the profile. The mean weight of Zn ranged between 88.26-201.65 mg/kg. According to the Igeo Index, the Zn concentrations of all soil samples fell into the uncontaminated category, except for the first horizons of Profiles 2 and 4, which were moderately contaminated (Table 6). The spatial interpolation of Zn is presented in Fig. 8. The highest concentration was in the south-west of the study area, which is located 270 m north of Mansuriyyat Al-Farastaq village. This could be due to atmospheric deposition, originating from local industrial plants. The highest Zn concentrations were in the swales top-soil and moderately high terrace units.




Fig. 8. Spatial interpolation of the weighted mean of zinc

Table 4. XRF analysis of soils collected from the study area

Profile no	Mapping unit	Depth (cm)	Metal concentrations (mg/kg)				
			V	Cr	Ni	Cu	Zn
1	T3	0-50	227.1	179.7	68.30	78.60	94.30
		50-85	265.9	158.1	77.80	60.90	85.40
		85-120	250.4	167.3	73.50	81.40	93.30
		120-150	221.8	163.3	61.60	60.10	80.80
2	S	0-45	744.4	519.0	267.30	288.90	377.60
		45-85	244.9	161.4	63.50	50.90	75.50
		85-110	250.8	192.4	70.20	71.40	86.80
3	D1	0-75	221.0	170.7	63.80	94.50	90.20
		75-100	241.4	179.6	63.00	73.00	86.90
		100-150	238.9	166.6	81.60	76.60	85.00
4	T2	0-60	203.1	159.2	72.30	118.70	308.00
		60-100	258.3	166.3	70.90	0.00	88.70
		100-120	222.8	166.2	65.50	75.20	84.90
		120-150	206.1	179.8	70.70	72.30	91.60
5	D2	0-45	194.0	149.1	74.40	95.20	103.10
		45-65	197.5	159.6	69.20	93.30	98.40
		65-110	225.3	143.0	73.20	95.50	95.00
		110-150	205.0	150.5	72.20	94.60	98.80
6	OB1	0-35	210.1	140.3	60.60	131.80	124.50
		35-65	216.0	152.7	76.70	76.40	109.40
		65-100	220.9	164.3	84.50	97.50	103.20
		100-150	219.5	153.7	85.00	0.00	100.60
7	OB2	0-55	228.2	180.8	76.50	93.50	94.70
		55-110	230.3	170.7	73.30	89.10	84.70
		110-150	196.5	168.7	64.50	74.10	84.30
8	L	0-30	218.7	164.1	68.90	0.00	93.20
		30-60	241.3	151.3	73.90	0.00	97.10
		60-100	247.3	155.4	82.10	0.00	89.50
		100-150	231.9	154.9	75.40	85.80	93.80
9	T1	0-45	250.7	156.8	78.20	0.00	112.10
		45-105	232.8	158.1	80.90	74.30	103.30
		105- 130	223.7	168.8	76.40	74.50	89.50
		130-150	231.1	168.4	69.70	74.10	94.00

Table 5. Heavy metal concentrations in soil samples and concentration limits recommended by Bowen (1979)

Profile No	Mapping unit	Mean weight of metals concentrations (mg/kg)				
		V	Cr	Ni	Cu	Zn
1	T3	240.53	168.48	70.39	71.42	89.29
2	S	450.58	314.73	148.39	152.92	201.65
3	D1	230.36	170.81	69.60	84.94	87.91
4	T2	221.04	166.14	70.70	71.96	176.49
5	D2	206.79	149.06	72.76	94.87	98.89
6	OB1	216.93	152.84	77.53	68.79	108.54
7	OB2	220.51	173.87	72.12	86.71	88.26
8	L	235.24	156.15	75.58	28.60	93.19
9	T1	236.42	160.86	83.28	52.01	102.40
Concentration limits (mg/kg)		90	70	50	25	90

Table 6. Igeo Index concentrations and associated contamination levels

Profile no	Depth (cm)	V	C Level	Cr	C Level	Ni	C Level	Cu	C Level	Zn	C Level
1	0-50	0.75	UN/M	0.77	UN/M	ND	UN	1.06	M	ND	UN
	50-85	0.97	UN/M	0.59	UN/M	0.05	UN/M	0.69	UN/M	ND	UN
	85-120	0.89	UN/M	0.67	UN/M	ND	UN	1.11	M	ND	UN
	120-150	0.71	UN/M	0.63	UN/M	ND	UN	0.68	UN/M	ND	UN
2	0-45	2.46	M/S	2.30	M/S	1.83	M	2.94	M/S	1.48	M
	45-85	0.85	UN/M	0.62	UN/M	ND	UN	0.44	UN/M	ND	UN
	85-110	0.89	UN/M	0.87	UN/M	ND	UN	0.92	UN/M	ND	UN
3	0-75	0.71	UN/M	0.70	UN/M	ND	UN	1.33	M	ND	UN
	75-100	0.83	UN/M	0.77	UN/M	ND	UN	0.96	UN/M	ND	UN
	100-150	0.82	UN/M	0.66	UN/M	0.12	UN/M	1.03	M	ND	UN
4	0-60	0.58	UN/M	0.60	UN/M	ND	UN	1.66	M	1.18	M
	60-100	0.93	UN/M	0.66	UN/M	ND	UN	ND	-	ND	UN
	100-120	0.72	UN/M	0.66	UN/M	ND	UN	1.00	M	ND	UN
	120-150	0.61	UN/M	0.77	UN/M	ND	UN	0.94	UN/M	ND	UN
5	0-45	0.52	UN/M	0.50	UN/M	ND	UN/M	1.34	M	ND	UN
	45-65	0.54	UN/M	0.60	UN/M	ND	UN	1.31	M	ND	UN
	65-110	0.73	UN/M	0.44	UN/M	ND	UN	1.34	M	ND	UN
	110-150	0.60	UN/M	0.51	UN/M	ND	UN	1.33	M	ND	UN
6	0-35	0.63	UN/M	0.41	UN/M	ND	UN	1.81	M	ND	UN
	35-65	0.67	UN/M	0.54	UN/M	0.03	UN/M	1.02	M	ND	UN
	65-100	0.71	UN/M	0.64	UN/M	0.17	UN/M	1.37	M	ND	UN
	100-150	0.70	UN/M	0.54	UN/M	0.18	UN/M	ND	-	ND	UN
7	0-55	0.75	UN/M	0.78	UN/M	0.02	UN/M	1.31	M	ND	UN
	55-110	0.77	UN/M	0.70	UN/M	ND	UN	1.24	M	ND	UN
	110-150	0.54	UN/M	0.68	UN/M	ND	UN	0.98	UN/M	ND	UN
8	0-30	0.69	UN/M	0.64	UN/M	ND	UN	ND	-	ND	UN
	30-60	0.83	UN/M	0.52	UN/M	ND	UN	ND	-	ND	UN
	60-100	0.87	UN/M	0.56	UN/M	0.13	UN/M	ND	-	ND	UN
	100-150	0.78	UN/M	0.56	UN/M	0.007	UN/M	1.19	M	ND	UN
9	0-45	0.89	UN/M	0.57	UN/M	0.06	UN/M	ND	--	ND	UN
	45-105	0.78	UN/M	0.59	UN/M	0.1	UN/M	0.98	UN/M	ND	UN
	105- 130	0.72	UN/M	0.68	UN/M	0.02	UN/M	0.99	UN/M	ND	UN
	130-150	0.77	UN/M	0.68	UN/M	ND	UN	0.98	UN/M	ND	UN

C Level = contamination level

UN = Uncontaminated

UN/M = Uncontaminated/moderately contaminated

M = Moderately contaminated

M/S = Moderately/strongly contaminated

ND = Not detected.

### 2.3. Major Soil oxides

Soil samples were analysed for heavy metals and major oxides (Table 7). Results for  $\text{SiO}_2$ ,  $\text{Al}_2\text{O}_3$ ,  $\text{P}_2\text{O}_5$ ,  $\text{K}_2\text{O}$ ,  $\text{CaO}$ ,  $\text{MgO}$ ,  $\text{Na}_2\text{O}$  and  $\text{Fe}_2\text{O}_3$  were compared with average concentrations of major oxides in soil (Bohn et al. 2001) (Table 5).  $\text{SiO}_2$  concentrations varied from 51.40-56.55% and were all less than the representative average value of 72.64% (Bohn et al. 2001).  $\text{Al}_2\text{O}_3$  concentrations varied from 16.28-24.42% and the mean concentration of profiles ranged from 18.81-22.9%.  $\text{Al}_2\text{O}_3$  concentrations in all samples exceeded the representative mean value of 13.22%.  $\text{K}_2\text{O}$  concentrations ranged from 1.05-1.62% with weighted mean values ranging from 1.14-1.35%, near the 1.2% representative mean.  $\text{CaO}$  concentrations ranged between 2.58-6.69% and the mean weighted value ranged between 3.70-5.95%.  $\text{CaO}$  and  $\text{Na}_2\text{O}$  concentrations exceeded the representative means of 1.44% and 0.99%, respectively.  $\text{Fe}_2\text{O}_3$  concentration ranged between 9.73-12.23%, whereas the representative mean is 5.77%.  $\text{P}_2\text{O}_5$  concentrations ranged from 0.15-0.49%. The weighted mean concentrations of  $\text{P}_2\text{O}_5$  samples exceed the 0.18% representative mean (Table 8). Thus, these deltaic soils are predominantly siliceous, with slight enrichment of the alumina component.



Table 7. Summary of major oxide concentrations in soil samples of the study area

Profile No.	Mapping unit	Depth (cm)	Major oxide concentrations (%)							
			SiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	CaO	MgO	Na <sub>2</sub> O	Fe <sub>2</sub> O <sub>3</sub>
1	T3	0-50	52.67	20.10	0.31	1.29	3.66	4.18	1.44	10.93
		50-85	56.46	23.92	0.25	1.29	3.18	4.53	1.44	12.03
		85-120	55.12	21.48	0.23	1.16	3.88	4.43	1.77	11.54
		120-150	54.42	19.28	0.21	1.19	4.17	3.98	2.02	10.38
2	S	0-45	ND	ND	ND	ND	ND	ND	ND	ND
		45-85	52.26	18.57	0.24	1.21	4.98	4.66	1.55	11.16
		85-110	55.35	21.50	0.20	1.10	4.50	4.62	1.89	11.28
3	D1	0-75	54.33	20.74	0.39	1.45	4.53	4.83	1.48	10.51
		75-100	52.83	18.36	0.20	1.10	5.03	4.30	1.72	11.20
		100-150	54.03	20.96	0.18	1.08	4.63	4.59	1.66	11.03
4	T2	0-60	55.87	19.11	0.43	1.60	3.70	5.11	1.49	10.61
		60-100	56.55	21.37	0.20	1.31	4.26	5.26	1.47	10.77
		100-120	53.60	18.24	ND	1.13	6.69	4.96	1.70	10.78
		120-150	52.15	16.48	0.20	1.07	6.34	4.55	1.84	10.28
5	D2	0-45	52.70	18.80	0.46	1.30	5.36	5.16	1.42	10.54
		45-65	54.20	19.19	0.20	1.24	4.38	5.43	1.43	10.96
		65-110	54.88	19.60	0.17	1.20	3.56	5.57	1.50	11.03
		110-150	53.93	19.20	0.28	1.25	4.43	5.39	1.45	10.84
6	OB1	0-35	54.64	19.19	0.49	1.62	5.22	4.99	1.48	10.82
		35-65	55.46	22.14	0.29	1.35	4.03	4.99	1.39	11.44
		65-100	55.71	23.59	0.20	1.25	3.47	5.00	1.36	11.74
		100-150	56.04	23.64	0.19	1.18	3.40	4.99	1.33	11.75
7	OB2	0-55	56.43	19.43	0.24	1.39	4.74	5.88	1.60	10.72
		55-110	55.29	19.4	0.15	1.22	3.06	4.99	2.44	10.59
		110-150	53.4	17.14	0.17	1.28	4.31	4.62	2.80	9.72
8	L	0-30	52.38	19.40	0.32	1.22	4.23	4.35	1.59	10.38
		30-60	54.42	19.47	0.28	1.21	4.35	4.90	1.62	11.11
		60-100	55.01	23.20	0.21	1.09	3.45	4.80	1.50	11.67
		100-150	56.06	24.42	0.17	1.09	2.58	4.79	1.40	12.23
9	T1	0-45	51.40	17.93	0.31	1.47	4.67	4.93	1.52	11.06
		45-105	52.32	19.01	0.20	1.14	4.32	4.78	1.64	11.43
		105- 130	52.36	21.47	0.16	1.05	3.12	4.52	1.70	11.73
		130-150	52.09	21.08	0.15	1.05	3.17	4.44	1.71	11.68

\*ND = Not detected.

Table 8. Mean weight of major oxide concentrations in soil samples and representative average limits (Bohn et al. 2001)

Profile No.	Mapping unit	Mean weight of major element concentrations (%)							
		SiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	CaO	MgO	Na <sub>2</sub> O	Fe <sub>2</sub> O <sub>3</sub>
1	T3	54.47	21.14	0.25	1.23	3.70	4.27	1.63	11.21
2	S	ND	ND	ND	ND	ND	ND	ND	ND
3	D1	53.97	20.40	0.28	1.27	4.64	4.66	1.58	10.79
4	T2	55.06	19.07	ND	1.35	4.77	5.01	1.57	10.60
5	D2	53.88	19.19	0.28	1.24	4.44	5.38	1.45	10.82
6	OB1	55.51	22.29	0.28	1.33	5.95	4.99	1.38	11.47
7	OB2	55.20	18.81	0.18	1.29	4.01	5.21	2.23	10.40
8	L	54.71	22.09	0.23	1.14	3.49	4.72	1.50	11.48
9	T1	52.01	19.36	0.22	1.21	4.07	4.73	1.62	11.40
Conc. limits (%)		70.29	>13.22	0.18	1.20	1.44	0.99	0.99	5.77

\*ND = Not detected.

## 2.4. Relationships between trace and major elements

V, Cr, Ni, Cu and Zn concentrations are significantly correlated (Table 9). There are no significant correlations between major elements and heavy metal concentrations, except for V, Ni and Zn. V and Ni have significant positive medium and strong correlations with Fe, respectively. This may indicate the sorption of these elements by Fe hydroxides. Ni has a strong positive association with Al, whereas a strong significant association was found between Zn and K<sub>2</sub>O and a medium correlation with P<sub>2</sub>O<sub>5</sub>. Such associations indicate strong affinity for these elements for Fe, Al and K oxides. Al<sub>2</sub>O<sub>3</sub> and Fe<sub>2</sub>O<sub>3</sub> display a strong positive significant correlation. In addition, P<sub>2</sub>O<sub>5</sub> is significantly correlated with K<sub>2</sub>O. There are strong significant negative correlations between Al<sub>2</sub>O<sub>3</sub> and CaO, between CaO and Fe<sub>2</sub>O<sub>3</sub> and a medium significant negative correlation between Na<sub>2</sub>O and Fe<sub>2</sub>O<sub>3</sub>.

Table 9. Correlation coefficients between trace and major elements in soils of the study area

	V	Cr	Ni	Cu	Zn	SiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	CaO	MgO	Na <sub>2</sub> O
Cr	0.97***											
Ni	0.97***	0.96***										
Cu	0.64***	0.71***	0.68***									
Zn	0.73***	0.75***	0.77***	0.68***								
SiO <sub>2</sub>	0.19	0.11	0.28	0.18	0.19							
Al <sub>2</sub> O <sub>3</sub>	0.4*	-0.13	0.56***	-0.17	-0.09	0.61***						
P <sub>2</sub> O <sub>5</sub>	-0.33	-0.35	-0.26	0.29	0.49**	-0.06	-0.22					
K <sub>2</sub> O	-0.25	-0.32	-0.19	0.32	0.56***	0.23	-0.18	0.78***				
CaO	-0.23	-0.15	-0.42**	0.14	-0.09	-0.38	-0.69***	0.37	0.1			
MgO	-0.25	0.35	0.27	0.24	0.21	0.41*	-0.03	0.14	0.39*	0.08		
Na <sub>2</sub> O	-0.1	0.40*	-0.33	-0.05	-0.20	-0.16	-0.43	-0.37	-0.21	0.06	-0.30	
Fe <sub>2</sub> O <sub>3</sub>	0.51**	-0.17	0.56***	-0.18	-0.11	0.26	0.79***	-0.31	-0.35	-0.55***	-0.07	-0.51***

Note: \*P &lt; 0.05, \*\*P &lt; 0.01, \*\*\*P &lt; 0.001, n = 33 soil samples.

## Conclusions

This study shows that concentrations of V, Ni and Cr exceeded recommended limits in the soils of the Middle Nile Delta. Cu concentrations were very variable. Zn concentrations slightly exceed the recommended limit. V, Cr, Ni, Cu and Zn concentrations are significantly correlated. There are no significant correlations between major elements and heavy metal concentrations, except for V, Ni and Zn. V and Ni have significant positive medium and strong correlations with Fe, respectively. The Igeo Index of V showed that all soil samples were in the uncontaminated/moderately contaminated categories, except for the first horizon of Profile 2, which is classified as moderately/strongly contaminated, while the Igeo Index for Cr showed most soil samples are in the uncontaminated/moderately contaminated category. The Igeo Index for Ni reveals that all samples are in the uncontaminated and moderately contaminated categories. For Cu, soil samples were in three contamination categories (uncontaminated/moderately contaminated, moderately contaminated and moderately/strongly contaminated). All Zn concentrations were in the uncontaminated category, except for the first horizons of Profiles 2 and 4, which were moderately contaminated. The highest heavy metal concentrations dominate the south-west of El-Gharbia Governorate and is mainly attributed to human activities, especially pollution from the Kfr Elzayat urban area (population in 1995 was 448,965). In terms of the distribution of heavy metals in the different physiographic units, the swales unit contained the highest values, as this is in Kfr El Zayat, which has many factories. We recommend that heavy metal contamination be studied within entire soil profiles and not just top-soils, because these metals affect soil and crop quality and can cause ground-water pollution. Protection against this hazard is vital for sustainable land management. Precise measures and efficient methods to improve soil and water quality must be conducted, in order to prevent soil and water pollution and to avoid the need for costly remediation in the future.

## References

- Abd Elgawad, M.; Hamdi, A. A.; Mahmoud, M. S.; Samir, I. G. 2007. Status of some heavy metals in Fayoum district soils: Third Conf. for Sustainable Agricultural Development, 12-14 Nov., Fayoum, Egypt. 507-526.
- Abdel-Moati, M. A. R.; El-Sammak, A. A. 1997. Man-made impact on the geochemistry of the Nile Delta Lakes. A study of metal concentrations in sediments. *Water, Air and Soil Pollution* 97(3): 413-429.
- Abdel-Sabour, M. F.; Abdou, F. M.; Elwan, I. M.; Al-Salama, Y. J. 2002. Effect of soil contamination due to wastewater irrigation on Cr fractions in some soils of Egypt. *Proc. 6th Radiation Physics Conf.*, Assuit, Egypt. 27-30.
- Abu Khatita, A. M. 2011. Assessment of soil and sediment contamination in the Middle Nile Delta area (Egypt)- Geo-Environmental study using combined sedimentological, geophysical and geochemical methods. German, Alexander-Universität Erlangen-Nürnberg zur Erlangung.
- Ali, A. A.; Moghanm, F. S. 2013. Variation of soil properties over the landforms around Idku Lake, Egypt. *The Egyptian Journal of Remote Sensing and Space Sciences* 16(1):91-101.
- Atwia, M. G.; Abou Heleika, M. M.; El-Shishtawy, A. M.; Sharp, J. M. 2006. Hydrostratigraphy of the Central Nile Delta, Egypt, using geoelectric measurements. *EGS-AGU-EUG Joint Assembly*, Nice, France 5:6-11.
- Baghdady, N. H.; Sippola, J. 1984. Extractability of polluting elements Cd, Cr, Ni and Pb of soil with three methods. *Acta Agriculturae Scandinavica* 34(3):345-348.
- Baker, D. E.; Senft, J. P. 1995. *Copper*. In: Alloway, B. J. (ed.): *Heavy Metals in Soils*. London: Blackie.
- Ball, D. F. 1964. Loss-on-ignition as estimate of organic matter and organic carbon in non-calcareous soils. *Journal of Soil Science* 15(1): 84-92.
- Bohn, L. H.; McNeal, L. B.; O'Connor, A. G. 2001. *Soil Chemistry*; 2nd edn, New York: John Wiley.
- Bowen, H. J. M. 1979. *Environmental Chemistry of the Elements*. London: Academic Press.
- Bradl, H.; Xenidis, A. 2005. Chemical remediation techniques, in Bradl, H. B. (Ed); *Heavy Metals in the Environment: Origin, Interaction and Remediation*. Amsterdam: Elsevier.
- Brough, P. A. 1986. *Principles of Geographical Information Systems for Land Resources Assessment*. Oxford: Oxford University Press.
- Climatologically Normal for Egypt. 2011. The normal for El-Gharbia Governorate Station (1960-2011), Ministry of Civil Aviation: Meteorological.
- Dawoud, M. A. 2004. Design of national groundwater quality monitoring network in Egypt. *J. Environ. Monit. Assess.* 96:99-118.
- Dobos, E.; Micheli, E.; Baumgardner, M. F.; Biehl, L. and Helt, T. 2000. Use of combined digital elevation models and satellite radiometric data for regional soil mapping. *Geoderma*. 97:367-391.
- Dobos, E.; Norman, B.; Bruce, W.; Luca, M.; Chris, J.; Erika, M. 2002. The use of DEM and satellite images for regional scale soil database. 17th World Congress of Soil Science (WCSS), 14-21 August, Bangkok, Thailand.
- Dumont, H. J. 2009. A description of the Nile Basin, and a synopsis of its history, ecology, biogeography, hydrology, and natural resources. In: Dumont, H. J. (Ed.) *The Nile: Origin, Environments, Limnology and Human Use*. New York: Springer.
- [http://link.springer.com/chapter/10.1007/978-1-4020-9726-3\\_1?no-](http://link.springer.com/chapter/10.1007/978-1-4020-9726-3_1?no-)
- Fanos, A. M. 2002. Report on the Modified Mega Delta Workshop. First Iteration 6 September.
- FAO 2006. *Guidelines for Soil Description*, 4th edn. FAO, Rome. ISBN 92-5-105521-1.
- Govil, P. K.; Reddy, G. L. N.; Krishne, A. K. 2001. Contamination of soil due to heavy metals in Patanchera Industrial Development Area: Andhra Pradesh, India. *Environ. Geol.* 41:461-469.
- Gowd, S. S.; Reddy, M. R. and Govil, P. K. (2010). Assessment of heavy metal contamination in soils at Jajmau (Kanpur) and Unnao industrial areas of the Ganga Plain, Uttar Pradesh, India. *J. Hazard. Mater.*, 174:113-121.
- He, J. Y.; Jia, X. 2004. Arc GIS geostatistical analyst application in assessment of MTBE contamination, ESRI User Conference Fremont, CA. Available at: <http://gis.esri.com/library/userconf/proc04/docs/pap1628>

- ITT. 2009. ITT Corporation ENVI 4.7 software, White Plains: New York.
- Jankaite, A.; Vasarevičius, S. 2005. Remediation technologies for soils contaminated with heavy metals. *Journal of Environmental Engineering and Landscape Management* 13(2):109-113.
- Lagerkvist J-S. B.; Oskarsson, A. 2007. Vanadium in Nordberg, G., Fowler, B., Nordberg, M. and Friberg, L., (Eds), *Handbook on the Toxicology of Metals*, Third ed. San Diego: Academic Press.
- Lillesand, T. M.; Kiefer, R. W. 2003. *Remote Sensing and Image Interpretation*. 2nd ed. New York: John Wiley.
- Lillesand, T. M.; Kiefer, R. W. 2007. *Remote Sensing and Image Interpretation*. 5th ed. New York: John Wiley.
- Ling, W.; Shen, Q.; Gao, Y.; Gu, X.; Yang, Z. 2007. Use of bentonite to control the release of copper from contaminated soils. *Australian Journal of Soil Research* 45(8):618-623.
- Lu, S. G.; Bai, S. Q. 2010. Contamination and potential mobility assessment of heavy metals in urban soils of Hangzhou, China. *Environ. Earth Sci.* 60(7):1481-1490.
- McLaughlin, M. J.; Zarcinas, B. A.; Stevens, D. P.; Cook, N. 2000. Soil testing for heavy metals. *Communications in Soil Science and Plant Analysis*, 31(11-14):1661-1700.
- Ohnesorge, F. K.; Wilhelm, M. 1991. Zinc in: E. Merian (Ed.), *Metals and their Compounds in the Environment, Occurrence, Analysis and Biological Relevance*. Verlagsgesellschaft, Weinheim, Basel: VCH.
- Rahman, S.; Khanam, D.; Adyel T.; Islam, M. Sh.; Mohammad Ahsan, A. and Akbor, M. A. (2012). Assessment of heavy metal contamination of agricultural soil around Dhaka Export Processing Zone (DEPZ), Bangladesh: Implications of seasonal variations and indices. *Appl. Sci.*, 2(3):584-601.
- Raven, P. H.; Berg, L. R.; Johnson, G. B. 1998. *Environment*, 2nd edition, New York: Saunders College Publishing.
- Reimann, C.; de Caritat, P. 1998. *Chemical Elements in the Environment*, Berlin: Springer-Verlag.
- Romic, M.; Romic, D. 2003. Heavy metal distribution in agricultural top-soils in urban areas. *Environmental Geology* 43:795-805.
- Said, R. 1993. *The River Nile: Geology, Hydrology and Utilization*. Oxford: Pergamon Press.
- Shepard, D., 1968. A Two-dimensional Interpolation Function for Irregularly-spaced Data. New York: ACM.
- Sunderman, F. W. (Jr.); Oskarsson, A. 1991. Nickel. In: Merian E (ed.) *Metals and their Compounds in the Environment; Occurrence, Analysis and Biological relevance*. Weinheim, Basel: VCH.
- Tong, S. T. Y. 1998. Indoor and outdoor household dust contamination in Cincinnati, Ohio, USA. *Environmental Geochemistry and Health* 20(3):123-133.
- USDA, 2010. *Keys to Soil Taxonomy*. United States Department of Agriculture, Natural Resources Conservation Service (NRCS) 11th Edition.
- WHO 1996. *Trace Elements in Human Nutrition and Health*. World Health Organization: Geneva.
- Wong, M. H.; Wong, J. W. C. 1986. Effects of fly ash on soil microbial activity. *Environ. Pollut., Ser A* 40:127-144.
- Woo, K. S.; Jo, J. H.; Basu, P. K.; Ahn, J. S. 2009. Stress intensity factor by p-adaptive refinement based on ordinary Kriging interpolation, *Finite Elem. Anal. Des.* 45(3):227-234.
- Zeydan, B. A. 2005. The Nile Delta in a Global Vision. Ninth International Water Technology Conference, IWTC9 2005, Sharm El-Sheikh, Egypt, 31-40.
- Zheng, C. 2006. Using multivariate analyses and GIS to identify pollutants and their spatial patterns in urban soils in Galway, Ireland. *Environmental Pollution*. 142(3):501-511.

**Mohamed S. Shokr:** Assistant Lecturer, received the degrees of B.Sc. and M.Sc. from Tanta University, Egypt. Currently, he is Assistant Lecturer in the Soils and Water Department, Tanta University and has a scholarship from the Egyptian Government to complete part of his Ph.D. studies in the University of Wolverhampton, UK. His research activities are concerned with pedology, soil sustainability, remote sensing and GIS.

**Ahmed A. El Baroudy:** Dr El Baroudy received the degrees of B.Sc., M.Sc. and Ph.D. from The University of Tanta, Egypt. Currently, he is Associate Professor of Soil in the Soil and Water Department of the University of Tanta. His research activities are mainly concerned with pedology, soil degradation, desertification, land evaluation, remote sensing and GIS modelling. He has published widely in Soil Science, having authored 21 refereed and conference papers. Dr El Baroudy has jointly supervised seven M.Sc. and five Ph.D. theses to completion. He has conducted postdoctoral research at Córdoba University, Spain. He is a member of the Editorial Board of the Comprehensive Research Journal of Agricultural Science (CRJAS). He also participates in five research projects. He is a member of many professional associations. These include the Egyptian Society of Soil Science, Journal of Agriculture Sciences of Tanta University, National Centre for Faculty and Leadership Development Egypt, International Centre for Faculty and Leadership Development at Tanta University. Dr El Baroudy is a manager of the Soil and Water Unit in the Faculty of Agriculture, Tanta University. He has won many awards, including the Tanta University Award for International Publishing, International Award for Agricultural Scientific Research in the Management and Development of Water Resources in Arid and Semi-arid Zones and the Tanta University Promotional Award in Agricultural Sciences.

Further information is available at the following web site:

[http://tdb.tanta.edu.eg/staff\\_data/Staff%20Detailed%20Data-ar.aspx?MemberID=3795](http://tdb.tanta.edu.eg/staff_data/Staff%20Detailed%20Data-ar.aspx?MemberID=3795).

**Michael A. Fullen:** Professor Fullen received the degrees of B.Sc. and M.Sc. from The University of Hull (UK), a Ph.D. from the UK Council for National Academic Awards (CNAA) and a D.Sc. from The University of Wolverhampton. Currently, he is Professor of Soil Technology at the University of Wolverhampton, UK. His research activities are mainly concerned with soil erosion, soil conservation, desertification and desert reclamation and his fieldwork is mainly based in Europe and Asia. He has published widely in Soil Science (as of March 2016, he has authored one book, 221 refereed papers, 217 conference papers and 26 consultancy reports). He is a referee for 48 journals and a member of the Editorial Board of 24 journals. Professor Fullen has jointly supervised 25 Ph.D. theses to completion and been Examiner for 25 Ph.D. theses. He has received research grants from over 30 institutions, principally the European Union. He is also Vice-President and UK Representative on the Council of the European Society for Soil

Conservation (ESSC) and Editor-in-Chief of the ESSC Newsletter. Professor Fullen is also WASWAC Vice-President, Councillor and Representative for the UK and was elected as an Academician of the Lithuanian Academy of Sciences. Professor Fullen was Joint Winner of the 'Gerold Richter Prize' from the ESSC in May 2015.

Further information is available at the following web site: [http://134.220.18.204/profiles/mike\\_fullen\\_cv.pdf](http://134.220.18.204/profiles/mike_fullen_cv.pdf)

**Talaat R. El-beshbeshy:** Professor El-beshbeshy received the degrees of B.Sc., M.Sc. and Ph.D. from The University of Minia, Egypt. Currently, he is Professor of Soil Science and Plant Nutrition in the Soil and Water Department of the University of Tanta. His research activities are mainly concerned with plant nutrition, hydroponics, soil fertility and soil chemistry. He has published widely in Soil Science, having authored 25 refereed and conference papers. He edited and translated one book (Principles of Plant Nutrition (in Arabic), Dar EL-Nashr for universities, Cairo, 1989). Professor El-beshbeshy has jointly supervised eight M.Sc. and 10 Ph.D. theses to completion. He has completed many scientific missions, Ph.D. graduate in Kassel University, Germany (1988-1990), Study visit to Kassel University, Germany (2 months during 1996), Visiting Professor at Kagawa University, Japan (3 months during 2001), Visiting Professor at Kassel University, Germany (2 months during 2005) and Visiting Professor at Kassel University, Germany (3 months during 2008). He is a member of many professional associations. These include the Soil Science Society of Egypt (SSSE), Egyptian Plant Society at Cairo University, Egyptian Society of Plant Nutrition and Fertilization at The National Research Center, Dokki, Cairo, Egypt, Journal of Agriculture Sciences, Tanta University, Deutschen Bodenkundlichen Gesellschaft and Alexandria Science Exchange at Alexandria University, Egypt.

**Rafat R. Ali:** Professor Ali has been Professor of Soil Science at the National Research Centre, Cairo, since 1999. He was appointed as a Research Assistant in the Soils and Water Use Department. In 1993, he was awarded the degree of M.Sc. and was then appointed as Assistant Researcher. He obtained a Ph.D. in 2013 and was then appointed as a Researcher. During 2003-2015 he published 56 scientific papers. He became Assistant Professor in 2008 and then Professor in 2014. He also participated in 24 research projects, published three book chapters and supervised several M.Sc. and Ph.D. degrees. He is a reviewer for seven international journals and assistant publisher of the three international journals. He received the 'Scientific Motivation in Agricultural Science' from the National Research Centre in 2008. In 2012 he received the first award of 'Excellence for Agricultural Scientific Research in Drylands' (ACSAD) from the League of Arab States. During 2004-2012, he worked as an expert in the National Authority for Remote Sensing and Space Sciences. Fields of interest include pedology, GIS, remote sensing and their applications in land degradation, land evaluation and landform mapping.

**Abd El Halim A. Abd El Halim:** Dr Abd El-Halim received the degrees of B.Sc. and M.Sc. from The University of Tanta and a Ph.D. from The University of Tanta through a scientific association with the University of Rostock, Germany. Currently, he is Associate Professor of Soil Physics and Water Relations at the University of Tanta. His research activities are mainly concerned with soil physics, soil erosion, soil conservation, irrigation water management, evapotranspiration models and desert reclamation. He has published widely in Soil Science (he has authored 12 refereed papers and three conference papers). He is a referee for four journals and a member of the Editorial Board of one journal. Dr Abd El-Halim has jointly supervised four M.Sc. and Ph.D. theses to completion. He has undertaken post-doctoral research at the Department of Soil Science, Kasetsart University, Nakhonpathom, Thailand.

**Antonio J.T Guerra:** Professor Guerra received the degrees of B.Sc. and M.Sc. from the Federal University of Rio de Janeiro (Brazil) and a Ph.D. from King's College London (UK). He undertook his first Post-doctoral Fellowship at Oxford University (UK) and his second at the University of Wolverhampton (UK). Currently, he is Professor of Physical Geography at the Department of Geography, Federal University of Rio de Janeiro, where he co-ordinates LAGESOLOS (the Laboratory of Environmental Geomorphology and Soil Degradation). His research activities are mainly concerned with soil erosion, land degradation, soil rehabilitation and his fieldwork is mainly based in Rio de Janeiro and São Paulo States (Brazil). He has published nearly 100 papers in refereed papers in both Brazilian and international journals. He has co-edited 17 books on geomorphology, soil erosion, environmental geomorphology, tourism, and soil rehabilitation in Brazil. Professor Guerra has supervised 21 Ph.D. theses and 25 M.Sc. dissertations to completion. He is a referee for 10 Brazilian and international journals. He has received research grants from over 15 Institutions. He is also Senior Researcher of the Brazilian Research Council (CNPq).

**Maria C.O. Jorge:** Mrs. Jorge received the degree of B.Sc. from the Federal University of Paraná (Brazil) and M.Sc. from the State University of Rio Claro (Brazil). Currently she is completing her Ph.D. at the Federal University of Rio de Janeiro and she has just finished the sandwich component of her Ph.D. at the University of Wolverhampton (2015). So far she has published 14 refereed papers in both Brazilian and international journals. She has also co-edited two books on land degradation and soil rehabilitation in Brazil. Her research activities are concerned with geotourism, geoconservation and geodiversity and her field work is conducted in Ubatuba Municipality, São Paulo State, Brazil.





## Journal of Environmental Engineering and Landscape Management

Saulėtekio al. 11, Vilnius-40, LT-10223, Lithuania

Email: [jeelm@vgtu.lt](mailto:jeelm@vgtu.lt)

Website: <http://www.tandf.co.uk/journals/TEEL>

### AUTHOR AGREEMENT

*This Agreement comes into effect if your Article is accepted for publication in the Journal of Environmental Engineering and Landscape Management (further refer as Journal)*

**Article title:** MAPPING OF HEAVY METAL CONTAMINATION IN ALLUVIAL SOILS OF THE MIDDLE NILE DELTA OF EGYPT (further refer as *Article*)

**Authors:** Mohamed S. Shokr, Ahmed A. El Baroudy, Michael A. Fullen, Talaat R. El-beshbeshy, Rafat R. Ali, Abd El Halim A. Abd El Halim, Antonio J.T Guerra and Maria C.O. Jorge.

**Corresponding author:** Mohamed Shokr (further refer as *Author*)

1. The Author hereby assigns to Vilnius Gediminas Technical University (the Publisher) the copyright in the Article and any supplemental tables, illustrations or other information submitted therewith that are intended for publication as part of or as a supplement to the Article in all forms, throughout the world, in all languages, for the full term of copyright. This assignment includes the right to provide the Article in electronic and online forms and systems.
2. The Publisher will permit the Author to use the Article elsewhere after publication, including placement it on the author's personal web pages, in other works or for the purposes of the Author's teaching and research, provided acknowledgement is given to the Journal as the original source of publication.
3. The Author warrants that the Article is original work, has not been published previously and is not under consideration for publication elsewhere. The Article contains no libellous or unlawful statements and that it in no way infringes the rights of others. The Author guarantees that the Author is the owner of the Article and has full power to make this agreement.
4. The Editor of the Journal is empowered to make such editorial changes as may be necessary to make the Article suitable for publication. The Editor will be made to consult the Author in any case if substantive changes are required.
5. If the Article was prepared jointly by more than one author, the Author warrants that he/she has been authorised by all co-authors to sign this agreement on their behalf.

Signed by the Author

Mohamed Shokr

Date 27/10/2015